

MULTIPLIERS FOR THE CONJUGATE SERIES OF A FOURIER SERIES

Prem Chandra

Date of Receiving : 06. 06. 2015

Date of Acceptance : 25. 06. 2015

Abstract. In this paper, we obtain multipliers for the absolute Riesz summability of the conjugate series of a Fourier series which not only improves some of the results obtained by R.Mohanty [Proc. London Math. Soc., 52(1951), 295-320] and P. Chandra [J. Orissa Math. Soc., 30(No. 2)(2011), 1-12] but also provides some new results.

1. Definitions and Notations

Let $\lambda(\omega)$ be the type and $\eta > 0$ be the order of Riesz mean. Then $\sum_{n=1}^{\infty} a_n$ is said to be summable absolutely by the type $\lambda(\omega)$ and order $\eta > 0$ of Riesz mean, symbolically we write,

$$\sum_{n=1}^{\infty} a_n \in |R, \lambda(\omega), \eta| (\eta > 0) \quad (1.1)$$

if ([7]; Definition B)

$$\int_h^{\infty} \{ \lambda^{(1)}(\omega) / \lambda^{(1+\eta)}(\omega) \} \left| \sum_{n < \omega} \{ \lambda(\omega) - \lambda(n) \}^{\eta-1} \lambda(n) a_n \right| d\omega < \infty, \quad (1.2)$$

where h is a suitable positive number ([9], [10]) and

$$\lambda^{(1)}(\omega) = \frac{d}{d\omega} \lambda(\omega). \quad (1.3)$$

Throughout, we use the following notations for a real fixed x and all real values of δ :

$$\psi(t) = \frac{1}{2} \{ f(x+t) - f(x-t) \} \quad (1.4)$$

$$\psi_1(t) = \frac{1}{t} \int_0^t \psi(u) du \quad (1.5)$$

$$\beta(t) = \psi(t) - \psi_1(t) \quad (1.6)$$

$$y_n^\delta = \log^{-\delta}(n+1) \quad (1.7)$$

$$b_n^\delta(t) = \int_t^\pi \log^\delta \frac{k}{u} d\left(\frac{\cos nu}{nu}\right) \quad (1.8)$$

2010 Mathematics Subject Classification. 42A45, 42A50, 40G99.

Key words and phrases. Conjugate series, Multipliers, Absolute Riesz summability.

Communicated by. Shashank Goel